




DEMOGRAPHICS

Total population (000)	191,972 (2008)
Total under-five population (000)	16,125 (2008)
Births (000)	3,105 (2008)
Birth registration (%)	89 (2005)
Under-five mortality rate (per 1000 live births)	22 (2008)
Infant mortality rate (per 1000 live births)	18 (2008)
Neonatal mortality rate (per 1000 live births)	11 (2008)
Total under-five deaths (000)	67 (2008)
Maternal mortality ratio (per 100,000 live births)	---
Lifetime risk of maternal death (1 in N)	---
Total maternal deaths	4,100 (2005)



Under-five mortality rate

Deaths per 1000 live births



Causes of under-five deaths, 2008

Globally more than one third of child deaths are attributable to undernutrition




MATERNAL AND NEWBORN HEALTH


Adolescent birth rate (births per 1,000 women)	56 (2006)
Unmet need for family planning (%)	7 (1996)
Antenatal visits for woman (4 or more visits, %)	89 (2006)
Intermittent preventive treatment for malaria (%)	NA*
C-section rate (total, urban, rural, %)	46, 35, 45 (2006)
Early initiation of breastfeeding (within 1 hr of birth, %)	43 (2006)
Postnatal visit for baby (within 2 days for home births, %)	---

Causes of maternal deaths

Regional estimates for Latin America / Caribbean, 1997-2007



Coverage along the continuum of care




INTERVENTION COVERAGE FOR MOTHERS, NEWBORNS AND CHILDREN

NUTRITION

Stunting prevalence (moderate and severe, %)	7 (2006)
Wasting prevalence (moderate and severe, %)	2 (2006)
Complementary feeding rate (6-9 months, %)	70 (2006)
Low birthweight incidence (%)	8 (2006)

Underweight prevalence


Percent children < 5 years underweight for age*



*Based on 2006 WHO reference population


Exclusive breastfeeding

Percent infants < 6 months exclusively breastfed




Vitamin A supplementation

Percent children 6-59 months receiving two doses of vitamin A during calendar year




Antenatal care

Percent women aged 15-49 years attended at least once by a skilled health provider during pregnancy




Skilled attendant at delivery

Percent live births attended by skilled health personnel



Neonatal tetanus protection


Percent of newborns protected against tetanus



CHILD HEALTH


Immunization

Percent of children immunised against measles
Percent of children immunised with 3 doses DPT
Percent of children immunised with 3 doses Hib




Diarrhoeal disease treatment

Percent children < 5 years with diarrhoea receiving oral rehydration therapy or increased fluids, with continued feeding




Malaria prevention

Percent children < 5 years sleeping under ITNs*




Malaria treatment

Percent febrile children < 5 years using antimalarials*




Prevention of mother to child transmission of HIV

Percent HIV+ pregnant women receiving ARVs for PMTCT



Pneumonia treatment


Percent children < 5 years with suspected pneumonia taken to appropriate health provider
Percent children < 5 years with suspected pneumonia receiving antibiotics



WATER AND SANITATION


Water

Percent population using improved drinking water sources



Sanitation


Percent population using improved sanitation facilities



EQUITY

Coverage gap by wealth quintile

Average of eight key indicators



POLICIES

International Code of Marketing of Breastmilk Substitutes	Yes
New ORS formula and zinc for management of diarrhoea	No
Community treatment of pneumonia with antibiotics	No
IMCI adapted to cover newborns 0-1 week of age	Yes
Costed implementation plan(s) for maternal, newborn and child health available	Partial
Midwives be authorised to administer a core set of life saving interventions	Partial
Maternity protection in accordance with ILO Convention 183	Partial
Specific notification of maternal deaths	Yes

SYSTEMS

Per capita total expenditure on health (US\$)	837 (2007)
General government expenditure on health as % of total government expenditure (%)	5 (2007)
Out-of-pocket expenditure as % of total expenditure on health (%)	34 (2007)
Density of health workers (per 10,000 population)	46.0 (2006)
Official Development Assistance to child health per child (US\$)	0 (2007)
Official Development Assistance to maternal and neonatal health per live birth (US\$)	0 (2007)
National availability of Emergency Obstetric Care services (% of recommended minimum)	---